

House of Learning COVID Return Flowchart

Children

What to do if a student is symptomatic at the Children's House of Learning

Symptoms of COVID-19

Not attributed to anther known condition

Fever (100.4) or chills cough

shortness of breath/difficulty

breathing

New loss of taste or smell

Muscle or body aches

Fatigue


Headache

Sore throat

Congestion or runny nose

Nausea or vomiting

Diarrhea



Quarantine: keeps someone who was in close contact with someone who has COVID-19 away from others. Quarantine if you have been in close contact with someone who has COVID-19, unless you have been fully vaccinated.

Isolation: keeps someone who is sick or tested positive for COVID-19 without symptoms away from others, even in their own home. People who are in isolation should stay home and stay in a specific "sick room" or area and use a separate bathroom (if available).

**3-ply disposable or N95 or KN95 mask

TOO-CDS 1/9/22

symptoms are

improving



House of Learning COVID Testing and Quarantine Guidance

HOUSE OF LEARNING EXPOSURE

Has this person received their booster or completed their vaccine series within the last 6 months (2 months for J&J)?



If symptom free, can continue at HOL uninterrupted. Test at 5 days after exposure

, No

Was the exposure during a low or high risk activity?

Low Risk Activity

May return to HOL immediately if no COVID symptoms. Test at 5 days from exposure to positive person.

High Risk Activity

*2 Years or older: May return to HOL immediately if no COVID symptoms and can mask regularly. Must be tested at days 2 and 4 from exposure. Quarantine for 5 days if no testing is done or mask cannot be worn regularly.

*When testing capability can accommodate. If limited supply, 2+ follow <2yr old guidance.

Younger than 2: Quarantine for 5 days, test at day 5 (If home test, follow testing instructions on test if 2 tests are needed).

Low Risk Activities: indoor classroom or extracurricular activities where masking is maintained. Exposed is within 3ft of an infectious person for 15+ mins.

High Risk Activities: Shared transportation, indoor meals, indoor unmasked sports, indoor classroom activity where masking is not maintained (students in kindergarten or younger, students with disabilities that preclude masking). Exposed is within 6ft of infections person for 15 or more minutes.

Quarantine: keeps someone who was in close contact with someone who has COVID-19 away from others. Quarantine if you have been in close contact with someone who has COVID-19, unless you have been fully vaccinated.

Isolation: keeps someone who is sick or tested positive for COVID-19 without symptoms away from others, even in their own home. People who are in isolation should stay home and stay in a specific "sick room" or area and use a separate bathroom (if available).

**3-ply disposable or N95 or KN95 mask


TOO-CDS 1/9/22



House of Learning COVID Testing Quarantine Guidance

HOUSEHOLD EXPOSURE

Has this person received their booster or completed their vaccine series within the last 6 months (2 months for J&J)?



Positive person able to isolate away from others?

Yes -> No quarantine. Test 5 days after last exposure to positive person. Continue to wear a**mask around others for 10 days.

If unable to mask, quarantine for 10 days.

Staff in toddler/infant rooms test every day at work from the first 5 days after exposure, as testing supplies allow.

No -> Quarantine for 5 days from positive person's symptom start date, test at day 5. Continue to wear a wear a mask** around other for 5 days after release from quarantine. If unable to mask, quarantine for 10 days.

Positive person is able to isolate away from others?

Yes -> Quarantine for 5 days from last exposure, test at day 5. Continue to wear a wear a mask** around other for 5 days after release from quarantine.

No -> Quarantine for 5 days from the recovery date of the positive person, **mask for 5 days after release from quarantine. If unable to mask, quarantine for 10 days.

Quarantine: keeps someone who was in close contact with someone who has COVID-19 away from others. Quarantine if you have been in close contact with someone who has COVID-19, unless you have been fully vaccinated.

Isolation: keeps someone who is sick or tested positive for COVID-19 without symptoms away from others, even in their own home. People who are in isolation should stay home and stay in a specific "sick room" or area and use a separate bathroom (if available).

**3-ply disposable or N95 or KN95 mask

TOO-CDS 1/9/22